

Advanced Calculus Entrance Exam

Spring 2001

- (1) Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be given by  $f(x, y) = (e^x \cos y, e^x \sin y)$ .
- (i) Show that  $Df(x, y)$  is invertible at every point of  $\mathbb{R}^2$ . 5%
  - (ii) Show that  $f$  is not one-to-one. 5%
  - (iii) Does (i) and (ii) contradict the *Inverse Function Theorem*? Why? 5%

- (2) Let  $R$  be a bounded closed set in  $\mathbb{R}^2$  and  $C$  be the smooth boundary curve. The *Green's second theorem* states:

$$\iint_R (u \Delta w - w \Delta u) dx dy = \int_C \left( u \frac{dw}{dn} - w \frac{du}{dn} \right) ds.$$

where  $w, u$  are both  $C^2$  functions on  $R$ .

- (i) How should you define the orientation of  $R, C$  and  $\vec{n}$  to make the formula correct? 5%
- (ii) How can you interpret this theorem as a formula for *Integration by parts*? 5%
- (iii) Let us define  $\langle f, g \rangle_R = \iint_R f(x, y) g(x, y) dx dy$  for any  $f, g \in \Omega$  where  $\Omega$  is the vector space of all  $C^\infty$  functions whose directional derivatives in the direction of normal at all points of  $C$  vanish. A linear operator  $L: \Omega \rightarrow \Omega$  is said to be *self-adjoint* if it satisfies  $\langle Lf, g \rangle_R = \langle f, Lg \rangle_R$ . Show that the Laplace Operator  $\Delta$  is self-adjoint. 5%

- (3) Let  $f$  be a continuous function of two variables  $(t, x)$  defined for  $t \geq a$  and  $x$  in some compact set  $S \subset \mathbb{R}$ . Assume that the integral

$$\int_a^\infty f(t, x) dt = \lim_{B \rightarrow \infty} \int_a^B f(t, x) dt$$

converges uniformly for  $x \in S$ .

- (i) Show that  $g(x) = \int_a^\infty f(t, x) dt$  is continuous for  $x \in S$ . 10%
  - (ii) Does  $\int_0^\infty x e^{-tx} dt$  converge uniformly for  $x \in [0, 1]$ ? Verify your answer. 10%
- (4) Let  $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$  be an invertible linear mapping and  $B_r$  be an  $n$ -dimensional ball centered at  $0$  with radius  $r$ . Compute 10%

$$\lim_{r \rightarrow \infty} \int_{T^{-1}(B_r)} e^{-\langle Ty, Ty \rangle} dy.$$

- (5) Let  $u = (u_1, u_2, \dots, u_n)^t \in \mathbb{R}^n$ ;  $f_j(u), j = 1, 2, \dots, q$  are continuously differentiable on  $\mathbb{R}^n$ . Consider 10%

$$L_p(u, \lambda) = \sum_{j=1}^q \lambda_j f_j(u) - (1/p) \sum_{j=1}^q \lambda_j \ln \lambda_j,$$

where  $p > 0$  and  $\lambda \in \Delta = \{ \lambda = (\lambda_1, \lambda_2, \dots, \lambda_q) \geq 0 \mid \sum_{j=1}^q \lambda_j = 1 \}$ . Show that, for each fixed  $p > 0$  and  $u \in \mathbb{R}^n$ , there is a unique optimal solution:

$$\lambda_j^*(u, p) = \exp(p f_j(u)) / \sum_{j=1}^q \exp(p f_j(u)), \quad j = 1, 2, \dots, q.$$

that maximizes  $L_p(u, \lambda)$  over  $\lambda \in \Delta$ .

- (6) Let  $l$  be a positive integer. Define

$$\Phi(\theta) = \frac{1}{l} \sum_{m=1}^l \cos \theta_m,$$

(背面仍有題目,請繼續作答)

where  $\theta = (\theta_1, \theta_2, \dots, \theta_l)$ . Further define

$$p(n, x, y) = \frac{1}{(2\pi)^l} \int_Q e^{i\langle \theta, (x-y) \rangle} \Phi^n(\theta) d\theta,$$

where  $x, y \in \mathbb{R}^l$ ,  $Q = \{\theta \mid -\pi \leq \theta_m \leq \pi, \forall m = 1, 2, \dots, l\}$  and  $\langle \theta, (x-y) \rangle$  is the usual inner product in  $\mathbb{R}^l$ . Consider

$$g(x, y) = \sum_{n=0}^{\infty} p(n, x, y).$$

- (i) Show that  $g(x, y) \leq \frac{1}{(2\pi)^l} \int_Q \frac{d\theta}{1 - |\Phi(\theta)|}$ . 10%
- (ii) Show that, there exists a neighborhood  $U$  of the point  $\theta = (0, 0, \dots, 0)$  in which 10%

$$\int_U \frac{d\theta}{1 - |\Phi(\theta)|} < \int_U \frac{4ld\theta}{\theta_1^2 + \theta_2^2 + \dots + \theta_l^2}.$$

(Hint: Use the first two terms of Taylor's expansion for each  $\cos \theta_j$ .)

- (iii) Use (ii) to show that  $g(x, y) < \infty$  for  $l \geq 3$ . (You may first try  $l = 3$  using Spherical Coordinates) 10%